


# Application of early error field correction to advanced scenarios

**J. Menard**

**NSTX FY2010 Research Forum  
LSB B318, PPPL  
December 2, 2009**



College W&M  
Colorado Sch Mines  
Columbia U  
CompX  
General Atomics  
INEL  
Johns Hopkins U  
LANL  
LLNL  
Lodestar  
MIT  
Nova Photonics  
New York U  
Old Dominion U  
ORNL  
PPPL  
PSI  
Princeton U  
Purdue U  
SNL  
Think Tank, Inc.  
UC Davis  
UC Irvine  
UCLA  
UCSD  
U Colorado  
U Illinois  
U Maryland  
U Rochester  
U Washington  
U Wisconsin

Culham Sci Ctr  
U St. Andrews  
York U  
Chubu U  
Fukui U  
Hiroshima U  
Hyogo U  
Kyoto U  
Kyushu U  
Kyushu Tokai U  
NIFS  
Niigata U  
U Tokyo  
JAEA  
Hebrew U  
Ioffe Inst  
RRC Kurchatov Inst  
TRINITI  
KBSI  
KAIST  
POSTECH  
ASIPP  
ENEA, Frascati  
CEA, Cadarache  
IPP, Jülich  
IPP, Garching  
ASCR, Czech Rep  
U Quebec

# Overview

## Brief Description:

- We propose to further optimize early  $n=1$  error-field correction for range of plasma scenarios and current ramp-rates
- **Such correction could be important for reduced density scenarios expected with the LLD**

## Background:

- In the FY2009 run, preliminary application of OH $\times$ TF  $n=1$  EFC was shown to increase early plasma rotation, reduce locking
- There are only 1 or 2 examples of such increased rotation and reduced locking (due to limited run time)
- More systematic investigation is warranted, since details of the EFC amplitude and turn-on time are important


# Early n=1 EFC using OH $\times$ TF EF compensation algorithm has significant impact on early plasma stability

- **Timing and amplitude scan for early OH $\times$ TF n=1 EF correction**

– Optimal correction


- EFC over-compensation can lead to  $\beta$  collapse

- Larger n=1 EFC over-compensation eliminates H-mode access




# Optimal early n=1 EFC reduces early locking tendency of n=1 tearing mode and substantially increases early rotation

- **Anti-corrective n=1 field (135778) vs. reference (135774)**




- *n=1 tearing mode nearly locks*




- *Rotation reduced 30-40%*
- *n=1 TM flattens rotation to low  $f_{\phi}$ =3-4kHz*

- **Optimal corrective n=1 field (135779) vs. reference (135774)**



- *n=1 tearing delayed, no locking, duration shortened*



- *Rotation increased 30%*
- *Core rotation maintained above 15kHz*



# Experimental Approach/Plan:

(1.0 day request, 0.5 day minimum useful)

- Reproduce increase in rotation with  $n=1$  early EFC
- Refine/scan EFC turn-on time, amplitude → optimize EFC to increase early rotation, reduce mode-locking activity
- Vary  $I_p$  ramp-rate (reduce it in steps of 1MA/s)
  - Vary EFC turn-on time and amplitude and assess impact
- Vary H-mode transition time during  $I_p$  ramp in steps of 20ms
  - Vary EFC turn-on time and amplitude and assess impact
- Determine EFC programming that performs well for widest range of cases, implement for subsequent discharges
  - Test on lower  $I_p$  and higher  $I_p$  case to confirm robustness

# Backup

---


# Motivation for “Early error-field correction in long-pulse plasmas”

- Insufficient fueling during LiTER generally results in “unstable” plasma early in discharge
    - Commonly attributed to “locked-modes”
      - Likely seeded by intrinsic error fields
    - But there are other effects of LiTER:
      - Confinement improvement from Li  $\rightarrow$  hit beta limit at fixed  $P_{\text{NBI}}$
      - Delayed H-mode mode, likely due to reduced density
    - Most (but not all!) EFC XPs rightly focused on sustaining high beta
  - Strong fueling during high-evap LiTER defeats purpose of Li
    - May not even be possible during (effective) LLD operation
- Reduced early EF could reduce mode locking, lower  $P_{\text{LH}}$ 
    - Now “know”  $n=3$  EF is from PF5  $\rightarrow$  early correction easy to test
    - $n=1$  EF caused by  $\text{OH} \times \text{TF}$ , and have correction algorithm in PCS

# 2006: XP614 demonstrated applying early n=1 EFC (based on OHxTF intrinsic EF) can increase early plasma rotation

Predictive OHxTF EFC on by t=150ms


EFC off



**Goal of XP-954 is to explore/extend these results further**




# Early n=3 EFC approx. proportional to PF5 current (known n=3 EF source) has modest impact on plasma evolution




**Early n=3 EFC induces earlier H-mode transition – reduced flow damping?**

# Both early n=3 EFC and NBI timing impact early H-mode transition

**Early n=3 EFC induces transition ~10ms earlier relative to late n=3 EFC**



**Rotation is ~10% higher with early n=3 EFC**



**Earlier NBI heating to 4MW also important for early transition**

# Early n=1 EFC using OH×TF EF compensation algorithm impacts H-mode access and confinement

- Compared to corrective polarity, anti-corrective polarity of n=1 field
  - reduces confinement,  $\beta$
  - delays early H-mode

