




NSTX



Supported by



U.S. DEPARTMENT OF ENERGY

Office of Science

# Study of chirping Toroidicity-induced Alfvén Eigenmodes in the National Spherical Torus Experiment


Mario Podestà

*R. E. Bell, A. Bortolon, N. A. Crocker, D. S. Darrow,  
E. D. Fredrickson, G.-Y. Fu, N. N. Gorelenkov,  
W. W. Heidbrink, G. Kramer, S. Kubota, B. P. LeBlanc,  
S. S. Medley, H. Yuh*

*and the NSTX Research Team*

**12<sup>th</sup> IAEA technical meeting on Energetic Particles in magnetic confinement systems**

**Austin, TX  
Sep. 7-10, 2011**




Columbia U  
CompX  
General Atomics  
FIU  
INL  
Johns Hopkins U  
LANL  
LLNL  
Lodestar  
MIT  
Nova Photonics  
New York U  
ORNL  
PPPL  
Princeton U  
Purdue U  
SNL  
Think Tank, Inc.  
UC Davis  
UC Irvine  
UCLA  
UCSD  
U Colorado  
U Illinois  
U Maryland  
U Rochester  
U Washington  
U Wisconsin

Culham Sci Ctr  
U St. Andrews  
York U  
Chubu U  
Fukui U  
Hiroshima U  
Hyogo U  
Kyoto U  
Kyushu U  
Kyushu Tokai U  
NIFS  
Niigata U  
U Tokyo  
JAEA  
Hebrew U  
Ioffe Inst  
RRC Kurchatov  
Inst  
TRINITI  
NFRI  
KAIST  
POSTECH  
ASIPP  
ENEA, Frascati  
CEA, Cadarache  
IPP, Jülich  
IPP, Garching  
ASCR, Czech Rep



# Bursting toroidicity-induced Alfvén eigenmodes (TAEs) can lead to enhanced fast-ion transport

- Multiple TAEs can be simultaneously destabilized
  - Possible overlap of many resonances in phase space
  - Non-linear development into “TAE *avalanches*”



# Bursting toroidicity-induced Alfvén eigenmodes (TAEs) can lead to enhanced fast-ion transport

- Multiple TAEs can be simultaneously destabilized
  - Possible overlap of many resonances in phase space
  - Non-linear development into “TAE *avalanches*” → fast ion losses



- Need to understand the physics of bursting TAEs, improve predictive capability for future devices (ITER)
- Spherical tokamaks such as NSTX provide excellent test-bed


# Outline

- Experimental scenario, diagnostics
- General features of TAEs on NSTX
  - Frequency, amplitude dynamics
  - Mode structure
  - Role of fast ion drive
- Non-linear dynamics, mode-mode coupling
- Summary and open issues

# Outline

- Experimental scenario, diagnostics
- General features of TAEs on NSTX
  - Frequency, amplitude dynamics
  - Mode structure
  - Role of fast ion drive
- Non-linear dynamics, mode-mode coupling
- Summary and open issues

# NSTX parameters




Major radius	0.85 m
<b>Aspect ratio</b>	1.3
Elongation	2.7
Triangularity	0.8
Plasma current	~1 MA
<b>Toroidal field</b>	<0.55 T
Pulse length	<2 s
<b>3 Neutral Beam sources</b>	
<b><math>P_{\text{NBI}} \leq 6 \text{ MW}</math>, <math>E_{\text{injection}} \leq 95 \text{ keV}</math></b>	
<b><math>1 &lt; v_{\text{fast}}/v_{\text{Alfvén}} &lt; 5</math></b>	

***This work:***  
***Focus on TAEs in L-mode plasma***  
***Center-stack limited***  
***Deuterium plasma***  
 ***$B_{\text{tor}}=0.55 \text{ T}$ ,  $I_p=0.7-0.9 \text{ MA}$***


# Mode activity and fast ion diagnostics on NSTX

shot#135404, t=320 ms



- Mirnov coils
    - Magnetic fluctuations up to 2.5 MHz
  - Multi-channel reflectometer
    - Mode structure (monotonic profiles)
- FFT analysis complemented by analysis in time domain to study mode dynamics over short time scale

UCLA




- Fast Ion diagnostics
  - Fast Ion D-Alpha systems (fast ion radial profile and spectrum)
  - Neutral Particle Analyzers, Fast Ion Loss probe, neutron rate

UCIrvine  
University of California, Irvine

# Experimental scenario :


$P_{NB} < 3\text{MW}$ ,  $n_e \sim 3-4 \times 10^{19}\text{m}^{-3}$ ,  $T_i \sim T_e = 1-1.5\text{keV}$



- NB-heated, L-mode plasmas
  - Plasma limited on center-stack
  - NB power and timing varied to affect mode stability
  - Plasma profiles evolving in time
    - > Monotonic, centrally peaked: OK for reflectometer measurements
  - Central plasma rotation up to 40kHz
    - > Large Doppler shift of mode frequency



# Safety factor profile evolves from strongly to slightly-reversed shear





- NB-heated, L-mode plasmas
  - Plasma limited on center-stack
  - NB power and timing varied to affect mode stability
  - Plasma profiles evolving in time
  - Reversed-shear  $q$  profile
    - >  $q_{\text{min}} \sim 1$  toward end of discharge
  - Safety factor evolution reconstructed through LRDFIT code constrained by MSE data

# Outline


- Experimental scenario, diagnostics
- **General features of TAEs on NSTX**
  - Frequency, amplitude dynamics
  - Mode structure
  - Role of fast ion drive
- Non-linear dynamics, mode-mode coupling
- Summary and open issues

# TAEs with low-intermediate toroidal mode number ( $n=2 \rightarrow 7$ ) are observed, with dominant $n=2-4$ modes



- Burst separation 0.5 – 2 ms
  - No systematic variation with  $n_e$ ,  $T_e$ ,  $P_{NB}$ , ...
  - Frequency evolution does not follow unique patten (e.g.  $t^{-1/2}$ , linear, exponential)
- Usually, each mode chirps independently of the others...
- ... but, eventually, *avalanches* occur:
  - Drop in neutron rate, FIDA

# Up to ~30% of fast ions can be lost during a single TAE avalanche



- Fast ion density (FIDA) drops over most of minor radius
- Loss results in a relaxation of the radial gradient → drive for TAEs is reduced
- Comparable losses estimated from FIDA and neutron rate
  - Large portion of phase space affected
- Losses increase with (total) mode amplitude
  - Linear? Quadratic?




# Frequency vs. amplitude are correlated; reminiscent of driven, non-linear system

- E.g., non-linear Van der Pol oscillator:

$$\ddot{x} - \epsilon(1 - \beta x^2)\omega_0 \dot{x} + (\omega_0^2 + \eta x^2)x = M\omega_0^2 \sin(\omega_D t)$$

damping      'non-linearity' factor      'restoring' term      driving force

- Can get info on damping, drive (resonant) frequency and their temporal variations?
- Comparison with chirping TAE data under way




# On average, TAE frequencies are consistent with a common frequency *in the plasma frame*

Measured frequency consistent with:

$$f_{lab,n}^{TAE} = f_0^{TAE} + n f_{Doppler}^{TAE}$$

↓ lab frame     ↓ plasma frame     ↓ shift from plasma rotation

- Valid for time scales >1 ms
- In general, each mode shows a different sub-millisecond dynamic




# On average, TAE frequencies are consistent with a common frequency *in the plasma frame*

Measured frequency consistent with:



$$f_{lab,n}^{TAE} = f_0^{TAE} + n f_{Doppler}^{TAE}$$

↓ lab frame     ↓ plasma frame     ↓ shift from plasma rotation

- Valid for time scales >1 ms
- In general, each mode shows a different sub-millisecond dynamic...
- ...except during large bursts:
  - Doppler shift only slightly changed here
  - Chirp mainly due to decrease in  $f_0^{TAE}$




FFT window 1.3ms



# Multi-channel reflectometer reveals details of mode structure and its temporal evolution

Crocker et al., PPCF 2011





- Makes it possible comparison with codes – e.g. NOVA – in terms of  $dn/n$



# NOVA eigenmodes selected based on measured frequency, mode number, structure






Fredrickson et al., PoP 2009

Fredrickson, P1.7





- NOVA provides multiple ideal eigenmodes in a given frequency range
- Match mode structure ( $dn/n$ ) from reflectometer to NOVA solutions
- Select 'observed' mode

# Mode structure maintain its shape even during strong, *multi-step* avalanches





- Outward propagation of *unstable front* during burst not observed Zonca et al., NF 2005
  - Broad mode structure, ~ minor radius
  - Incomplete transition TAE → EPM?





# Modes have broad mode structure, extending over good fraction of minor radius; peak at mid-radius



- sh#141711
- t=470ms
- Magnetic axis:  
R~105 cm
- Edge: R~155 cm








# Comparison with NOVA (ideal MHD) is satisfactory; however, no good match for $n=1$ 'TAE'



- Frequency from NOVA consistent with experiment, once Doppler shift is included
- Shift of 2-5 cm between measured and simulated  $dn/n$


- NOVA results assume ideal MHD, no fast ion effects
- *Need to carefully check profile consistency between experimental data (MPTS, CHERS) and TRANSP output / NOVA input*

# Initial results from [non-linear] self-consistent code M3D-K agree with experiments, too



- NOVA results assume ideal MHD, no fast ion effects
- *Need to carefully check profile consistency between experimental data (MPTS, CHERS) and TRANSP output / NOVA input ... and M3D-K*



# Preliminary results from M3D-K also show little effects of rotation on TAE structure



- No fast ion effects included
- Scale rotation profile from measured  $f_0=36$  kHz down to  $f_0=18$  kHz (50% reduction)
- Mode peak location, structure do not change substantially

*Fu, TTF 2011*


# Similar features observed in L- and H-mode plasmas and during combined NB+RF : robust dynamics



- Example: H-mode, NB / NB+RF
  - Profiles different from L-mode
  - Higher safety factor than for L-mode discharges
  - Reversed shear in both L- and H-mode

Fredrickson, P1.7


# Onset of bursting/chirping regime strongly dependent on injected NB (and RF) power



- Each symbol represents values for single  $n$  mode, 5ms average
  - $f_{0,pl}$ : mode frequency in the plasma frame




# High ratio of fast ion to thermal $\beta$ leads to bursts/chirps – but other factors determine mode dynamics




- Modes show more bursting character as discharge evolves
  - NB power increases, fast ion population and  $\beta_{f,i}$  builds up
  - Typical bursts have *edge*  $\delta B/B \sim 10^{-4}$  (from Mirnov coils)
    - > Can increase more than x10 during avalanches

# Destabilization of TAEs correlates with variations of heating scheme, e.g. NB vs. NB+RF




- Example: H-mode discharges with NB and NB+RF heating
  - Different profiles with respect to L-mode (e.g. higher safety factor)
  - Reversed shear in both L- and H-mode



**NB only**  
**NB+RF, NB notch**  
**NB+RF, RF notch**


- Here NB alone does not drive TAEs
- NB+RF does: increase in fast ions

# Fast ion population is key parameter to destabilize TAEs; subtle details hard to get experimentally...




**NB only**  
**NB+RF, NB notch**  
**NB+RF, RF notch**

- Marginal variations in  $n_f(R)$ ,  $\text{grad}(n_f)$  from FIDA, except for 100% change in source
- Can not differentiate between role of  $\beta_{f.i.}$  and radial gradient
- OK to look at trends



# Strong dependence on NB drive leads to entangled evolution of TAEs, fast ions and thermal plasma



- Mode location,  $R^{TAE}$ , obtained by matching  $f_{Doppler}^{TAE}$  with measured rotation profile:

$$f_{rot}(R^{TAE}) = f_{Doppler}^{TAE}$$

- Correlation between

- Mode location  $R @ f_{Doppler}^{TAE}$
- Max fast ion gradient  $max grad(n_f)$
- Max rotation shear



Coupling through common  
“source term”, i.e. **NB injection**



Modes cluster at similar radius:  
may enhance coupling

Podestà et al., PoP 2010

# Once set up, bursting/chirping TAE regime is rather insensitive to variations in plasma parameters





- Burst separation 0.5 – 2 ms
  - No systematic variation with  $n_e$ ,  $T_e$ ,  $P_{NB}$ , ...
  - Frequency evolution does not follow unique patten (e.g.  $t^{1/2}$ , linear, exponential)
- Usually, each mode chirps independently of the others...
- ... but, eventually, *avalanches* occur:
  - **Multiple modes lock on similar dynamic, mode-mode coupling**

# Outline

- Experimental scenario, diagnostics
- General features of TAEs on NSTX
  - Frequency, amplitude dynamics
  - Mode structure
  - Role of fast ion drive
- **Non-linear dynamics, mode-mode coupling**
- Summary and open issues

# Coupling between multiple TAEs, enhanced losses observed during explosive modes' growth




- Coupling generates higher/lower frequency modes

- Multiple modes follow similar dynamic during the burst
  - *Transition* from single- to multi-mode regime

Podestà et al., NF 2011

# 'New' modes appear in the spectrum above/ below TAE range during large bursts

- Modes can be classified into three groups
  - Discriminant: frequency, temporal evolution*



- Picture consistent with primary TAEs
  - coupling to each other
  - generating/pumping *secondary* modes through sum/difference with  $\Delta n=1$



# Simple model based on quadratic interactions can be used to investigate coupling between TAEs

$$\dot{s}_{n_3} = \langle c_{(n_1, n_2)} s_{n_1} s_{n_2} \rangle_{f_{n_3}}$$

$s_{n_2} \rightarrow s_{n_2}^*$  (complex conjugate) for difference interaction

Right-hand side filtered around frequency  $f_{n_3}$




Modes must satisfy matching conditions  $\begin{cases} n_3 = n_1 \pm n_2 \\ f_{n_3} = f_{n_1} \pm f_{n_2} \end{cases}$

$c(n_1, n_2)$  is the coupling coefficient, here used as 'free scaling parameter'

In practice:





- Real signals  $s_{n_1}$ ,  $s_{n_2}$ ,  $s_{n_3}$  measured for each possible triplet, e.g. from Mirnov coils
- "Reconstruct"  $\dot{s}_{n_3} \rightarrow \dot{s}_{n_3, rec}$  from measured  $s_{n_1}$ ,  $s_{n_2}$
- Compare measured and reconstructed  $\dot{s}_{n_3}$
- Frequency match must be verified in the plasma frame:
  - Rotation profile and location of each mode must be accurately known

# Good agreement with quadratic interactions' model: amplitude evolution and frequency matching



- $n=1$  mode fades away when either amplitude of pump modes or frequency matching vanishes


# Mode number matching condition verified



- Symbols: rms mode amplitude data from 11 Mirnov coils
- Solid lines: fit for a given  $n$  ( $n=1$  here)
- Dashed line: unit circle (zero-amplitude reference)


- “Reconstructed” toroidal structure of  $n=1$  mode also agrees with measured one
  - Phase shift of  $\sim 180$  degrees during strongest mode activity

# Phase matching condition is *transiently* verified for tens of (primary) mode cycles during large bursts




- Stationary phase during quadratic interaction is important!
  - $n=1$  mode fades away  $\Leftrightarrow$  phase changes rapidly in time
  - “Single mode” dynamic, with each mode following its own chirp/burst cycle, is effective in reducing efficiency of quadratic interactions
  - Result: small bursts (single mode), and occasional multi-mode avalanches
- $n=1$  mode mediates coupling: what is it?

# $f_{n=1}$ consistent with central plasma rotation: avalanches drive $f_0^{TAE} \rightarrow 0$ (plasma frame), cause coupling to 'kinks'?




- Kink-like modes destabilized during avalanche
- Sustained fast ion losses
- Reconstructed  $f_0^{TAE} \rightarrow 0$
- TAE mode structure  $\sim$  maintained
- Doppler shift radius  $R^{TAE}$  sweeps from plasma center out in  $\sim 2$  ms

Bortolon, O-12




n=1 kink-like + harmonics





# Meaning of $f_0^{TAE}$ : mode frequency at the mode location $R^{TAE}$ , i.e. where drive is maximum

TAE continuum from NOVA-K





# Meaning of $f_0^{TAE} \rightarrow 0$ : $f^{TAE}$ beats with $n \times$ rotation frequency at the plasma center

TAE continuum from NOVA-K



- Coupling with kink-like modes favored when  $n_{tor} \times f_{rot}$  on axis  $\sim f^{TAE}$
- Observed for  $q > 1$ : fishbone branch involved?



# $n=1, m=-2, \dots, +2$ kink found through ideal MHD code NOVA with no rotation, free boundaries



- Large edge perturbation, consistent with reflectometer's data





# Measured perturbation evolves during burst; large edge component, $dn/n > 5\%$



- Data from inversion of multi-channel reflectometer signals
- Assume 'linear' model for reflectometer, pure O-mode propagation, fixed  $n_{0,eI}$
- Roughly consistent with (ideal) NOVA solution

# MHD unstable $n=1$ kink also found through self-consistent M3D-K code



## **M3D-K:**

- *Realistic geometry (spherical tokamak)*
- *Experimental parameters and profiles (interface with TRANSP)*
- *Global; linear/nonlinear*
- *Non-perturbative fast ion effects (i.e., can model EPM) – not used here*
- *Plasma rotation is included*

**Fu, TTF 2011**

- Structure is different between NOVA and M3D-K
- Support hypothesis of marginally stable 'kink', destabilized during avalanche

# Outline

- Experimental scenario, diagnostics
- General features of TAEs on NSTX
  - Frequency, amplitude dynamics
  - Mode structure
  - Role of fast ion drive
- Non-linear dynamics, mode-mode coupling
- Summary and open issues

# Summary

- After onset, bursting/chirping TAE regime is “robust” against small variations of plasma parameters
  - L-mode vs. H-mode, NB only, NB+RF, ... : all show similar features
- Non-linearities at play in both single-mode and multi-mode (avalanching) TAE dynamics
  - But only avalanches seem to cause significant fast ion losses
  - Coupling manifests as intermittent, chaotic process
  - Coupling can encompass multiple ‘scales’: TAE, kink/fishbones (, GAE, ...)
  - Dynamics complicated by link between fast ion, TAEs, thermal plasma through NB injection
- Different physics for weak chirps vs. avalanches?
  - Weak chirps regulated by phase-space, fast ion profile effects; single-mode
  - Avalanches lock on underlying kink/fishbone (‘global’); multi-mode
- Present experiments allow thorough benchmark of codes
  - Linear MHD satisfactory for first-order estimates of mode structure
  - Non-linear, self consistent codes required to capture full dynamics

# Open Issues

- How to scale present experiments to future devices (ITER), burning plasmas?
  - Code validation, multi-machine comparison are the key
    - > Linear codes (NOVA)
    - > Quasi-linear models, self-consistent codes (M3D-K)
    - > Investigate wave-fast ion interaction (ORBIT, SPIRAL)
    - > *Improve stability calculations*
  - But: conditions will be very different, e.g.  $F_{f,i}(\underline{r},\underline{v},t)$  from alpha's (+NB, RF, ...)
    - > E.g. need better understanding of role of spatial vs. velocity gradients
  - Parallel effort: develop control tools
    - > Act on NB deposition, q-profile? Active antennas?
- Combine different diagnostics/techniques; develop phase-space models to interpret results consistently


# Backup slides

# Experimental scenario :



$P_{NB} < 3\text{MW}$ ,  $n_e \sim 3-4 \times 10^{19} \text{m}^{-3}$ ,  $T_i \sim T_e = 1-1.5 \text{keV}$

## NB-heated, L-mode plasmas

- Plasma limited on center-stack
- NB power and timing varied to affect mode stability
- Plasma profiles evolving in time
  - > Monotonic, centrally peaked: OK for reflectometer measurements
- Central plasma rotation up to 40kHz
  - > Large Doppler shift of mode frequency
- Reversed-shear  $q$  profile
  - >  $q_{\min} \sim 1$  toward end of discharge
- Safety factor evolution reconstructed through LRDFIT code constrained by MSE data






# No detectable fast ion losses are observed during weakly bursting/chirping phase



- Statistical average over ~20 events (~10ms)
- No clear evidence of losses from neutrons, FIDA
  - Does not exclude “continuous” (non-bursting) losses




# Up to ~30% of fast ions can be lost during a single TAE avalanche




- Fast ion density (FIDA) drops over most of minor radius
- Loss results in a relaxation of the radial gradient → drive for TAEs is reduced
- Comparable losses estimated from FIDA and neutron rate
  - Large portion of phase space affected
- Losses increase with (total) mode amplitude
  - Linear? Quadratic?

# Fast ion loss probe at vessel wall captures transient loss, provides details of affected phase space region

## Scintillator probe, sFLIP:




- ‘Magnetic spectrometer’
- Provides  $\Gamma_{\text{loss}}(\rho, \chi, t)$
- $5\text{cm} \leq r \leq 60\text{cm}$
- $15^\circ \leq \chi \leq 80^\circ$
- $f_{\text{sampl}} = 30\text{ kHz}$



- Observed losses at  $E \sim E_{\text{inj}}$ , wide pitch range
  - Possible signature of phase space stochastization
- Phase space model help understand loss process, measurements **Darrow, O-1**
- Simulations with ORBIT code under way

# Evidence of redistribution at play – multiple frequency scales can be involved



- Bursting GAEs sometimes observed at  $f \sim 1 \text{ MHz}$  during TAE activity
- GAE bursts precede TAE avalanches
  - But not all TAE avalanches are preceded by GAE burst
- Indication of fast ion redistribution by GAEs?
  - Different mode localization with respect to TAEs
- Losses only observed after TAE avalanches

*Fredrickson, IAEA 2010*



# M3D-K – non-linear, self-consistent Hybrid/MHD code

G.-Y. Fu et al., Phys. Plasmas 13, 052517 (2006)


*M3D-K code developed at PPPL and used to investigate fast-ion driven Alfvénic modes and MHD*

- 3-D nonlinear.
- Several different physical models:
  - Resistive MHD.
  - Hybrid (fluid electrons, particle ions).
  - MHD/particle (one-fluid thermal plasma, + energetic particle ions).
- Full-orbit kinetic ions.
- Drift-kinetic electrons.
- For particles: Drift-kinetic or gyrokinetic.

*M3D-K simulation of single vs. multi-mode TAE dynamics*




# NSTX reflectometers are used to measure density fluctuations



NSTX Multi-channel reflectometer system (UCLA)

- N. Crocker et al., PPCF 2011 (in press)





- Low-field side launchers/receivers
- Assume pure O-mode propagation

$$\Delta\Phi = \frac{2\omega}{c} \int_{x_{c.o.}}^{x_{inj}} N dx$$

$$N^2 = \frac{k^2 c^2}{\omega^2} = \frac{c^2}{\omega^2} k_{inj}^2 \left[ 1 - \frac{n(x)}{n_{c.o.}} \right]$$

# Analysis of reflectometer's data




- Low-field side launchers/receivers
- Assume pure O-mode propagation

$$\Delta\Phi = \frac{2\omega}{c} \int_{x_{c.o.}}^{x_{inj}} N dx$$

$$N^2 = \frac{k^2 c^2}{\omega^2} = \frac{c^2}{\omega^2} k_{inj}^2 \left[ 1 - \frac{n(x)}{n_{c.o.}} \right]$$

- Invert data by using *response matrix*,  $M : \Delta\Phi_i = M_{ij} \delta n_j$
- Find total  $\delta n$ , *then* filter around each mode's frequency
- OK for small ( $<3\%$ )  $\delta n$
- *Assumptions (pure O-mode, linear, ...) likely to break down for large perturbations*


# NOVA analysis reveals significant contribution to density perturbation from compressional term



- Compressional term is significant
- Impact interpretation of reflectometer's data, etc.

$$\frac{\delta n}{n} = \nabla \underline{\xi} - \underline{\xi} \cdot \frac{\nabla n}{n}$$

# Multi-channel reflectometer is used to look for decorrelation of TAE across rotation shear layer






## Procedure:

- Band-pass filter signals around mode frequency (ex.  $n=3$ )
- Reconstruct frequency, amplitude evolution
- Get cross-correlation
- Obtain cross-phase between channels from time delay



# Results from reflectometer show no evidence for spatial decorrelation of the modes across shear layer


- No difference in measured frequency
- Mode starts decaying when cross-correlation is still  $\sim 1$
- No systematic cross-phase variation during chirps




Podestà et al., PoP 2010

# No temporal correlation is observed between rotation shear evolution and mode amplitude rise/decay rates


- Effective rise/decay rates calculated from exponential fit (B-dot data)




- Shear increases in time
- No correlation with inferred rates





# Statistically significant ensemble does not show any clear trend of rise/decay rates on rotation shear



- Database of 10 discharges
- Group data with similar  $n_e$ ,  $T_e$ 
  - Similar damping on thermal plasma
- No simple dependence on injected NB power,  $T_e$ ,  $n_e$ , ...
- Trends hidden by changes in  $q$  profile, gap structure, ... ?




# Additional RF power marginally successful in decoupling fast ion, TAE and thermal plasma dynamics



- May explain slight differences between NB-only and NB+RF


# Bicoherence suggests coupling between modes at play during large bursts

- High bicoherence  $>70\%$  measured during burst
  - Average over 11 Mirnov coils distributed toroidally over  $360^\circ$
  - Indicative of sum/difference interactions between modes
  - Both TAEs and low-frequency modes participate




# Proximity of beam ions to sFLIP detector at high $\chi$ (i.e. high $\mu$ ) indicates why loss appears there first

shot#: 141707a shotTime: 485




## Solid State NPA diagnostic on NSTX operated in “current mode” during FY-10 for high temporal resolution data on mode-induced fast ion losses



- Measurement of flux of energetic neutrals
- 4 lines of sight on the NBI
  - $R_{\text{tan}} = 60, 90, 100, 120 \text{ cm}$
- Pinhole & Silicon photodiode detector (AXUV)
- Aluminum foil (150nm) blocks light, SXR, low energy neutrals (<10 keV)
- Detected neutrals generated by Charge Exchange of fast ions with **beam** and/or **edge** neutrals

# 'Fishbone beacon' observed during transient fast-ion losses associated with chirping modes



- Fast frequency chirping instabilities (e.g. TAE) accompanied by periodic bursts in ssNPA signals
- Fast ion loss cone, rotating in phase with the mode, is inferred based on time delay between ssNPA channels
- **Burst appear at different phases in different channels**