

# Progress in characterization of the pedestal structure, stability and fluctuations during ELM cycle on NSTX

**Ahmed Diallo<sup>1</sup>**

J. Canik<sup>2</sup>, T. Goerler<sup>3</sup>, G.J. Kramer<sup>1</sup>, S-H. Ku<sup>1</sup>, J. Manickam<sup>1</sup>, R. Maingi<sup>2</sup>,  
 D. Smith<sup>4</sup>, P. Snyder<sup>5</sup>, T. Osborne<sup>5</sup>, R.E. Bell<sup>1</sup>, D. Boyle<sup>1</sup>, C-S. Chang<sup>1</sup>,  
 W. Guttenfelder<sup>1</sup>, B.P. LeBlanc<sup>1</sup> and S. Sabbagh<sup>6</sup>

<sup>1</sup>Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ, USA

<sup>2</sup>Oak Ridge National Laboratory, Oak Ridge, TN, USA


<sup>3</sup>Max-Planck-Institut für Plasmaphysik-IPP, Garching, Germany

<sup>4</sup>University of Wisconsin, Madison, WI, USA

<sup>5</sup>General Atomics, San Diego, CA, USA

<sup>6</sup>Columbia University, New York, NY, USA

**APS-DPP 2012**  
**31st Oct Nov 2012**  
**Providence, RI**



*Culham Sci Ctr  
 York U  
 Chubu U  
 Fukui U  
 Hiroshima U  
 Hyogo U  
 Kyoto U  
 Kyushu U  
 Kyushu Tokai U  
 NIFS  
 Niigata U  
 U Tokyo  
 JAEA  
 Inst for Nucl Res, Kiev  
 Ioffe Inst  
 TRINITI  
 Chonbuk Natl U  
 NFRI  
 KAIST  
 POSTECH  
 Seoul Natl U  
 ASIPP  
 CIEMAT  
 FOM Inst DIFFER  
 ENEA, Frascati  
 CEA, Cadarache  
 IPP, Jülich  
 IPP, Garching  
 ASCR, Czech Rep*




*Coll of Wm & Mary  
 Columbia U  
 CompX  
 General Atomics  
 FIU  
 INL  
 Johns Hopkins U  
 LANL  
 LLNL  
 Lodestar  
 MIT  
 Lehigh U  
 Nova Photonics  
 Old Dominion  
 ORNL  
 PPPL  
 Princeton U  
 Purdue U  
 SNL  
 Think Tank, Inc.  
 UC Davis  
 UC Irvine  
 UCLA  
 UCSD  
 U Colorado  
 U Illinois  
 U Maryland  
 U Rochester  
 U Tennessee  
 U Tulsa  
 U Washington  
 U Wisconsin  
 X Science LLC*

# Abstract

Understanding the pedestal structure is important for achieving high performance pedestals necessary for maximum core fusion gain in ITER and future next-step devices. The stability of the pedestal is characterized in high performance discharges in National Spherical Torus Experiment (NSTX). In addition, the spatial structure of turbulence present during an ELM cycle in the pedestal region indicates spatial scales  $k_{\perp} \rho_i^{\text{ped}}$  ranging from 0.2 to 0.7 propagating in the ion diamagnetic drift direction at the pedestal top. These propagating spatial scales are found to be poloidally elongated and consistent with ion-scale microturbulence. Linear gyrokinetic simulations using GENE indicate the presence of hybrid ITG/KBM-TEM modes at the pedestal top. Nonlinear gyrokinetic simulations -- XGC1 -- find localized fluctuations agreeing with experimental level radial and poloidal correlation lengths.

*This work is supported by U.S. Dept of Energy contracts DE-AC02-09CH11466.*

# Radial profiles of density, temperature and velocity are composite of times between multiple fraction of ELMs (e.g., 50-99% ELM cycle)



Electrons edge radial profiles are fitted using a modified tanh function


$$N(\psi) = A \tanh \left( \frac{\psi_{sym} - \psi}{\psi_{width}} \right) + offset$$

R. Groebner and T. Osborne PoP (1998)




Ion profiles are spline fitted

# Temperature pedestal height increases during the ELM cycle while the density pedestal shows no convincing trend

- More than a factor of two increase in pedestal temperature
- Density pedestal is much less sensitive to the ELM cycle
- Heat and particle evolutions appear to be decoupled









# Pedestal width and height progressively increase during ELM cycle but the peak pressure gradient remains clamped



- Pedestal height increases by a factor  $\leq 3$ 
  - Height scales with  $I_p$
- Pedestal width increases independently of  $I_p$
- Gradient is clamped early in ELM cycle



# Ideal ballooning stability is performed using “ball” module of GS2 during the last 50% of ELM cycle: Pedestal top is found to be ballooning unstable



See Canik - this meeting

# ELITE Peeling-ballooning mode stability

## diagram confirms that NSTX pedestal is kink-peeling unstable



C-S Chang Wednesday P07.007

Equilibria around experimental point are generated by varying the edge pressure gradient at fixed edge current and vice-versa.

Stability of each equilibrium is computed using the ELITE MHD code


–  $n = 3, 6, 12, 15$

Using the recent implementation of XGC0-bootstrap current model, the experimental point is in the kink/peeling unstable region during the last part of the ELM cycle.

- The XGC0 current model is described in C-SChang TH/P4-12
- These results agree with previous NSTX stability analyses.

Boyle PPCF (2011)

# NSTX measured pedestal pressure width scales like $(\beta_\theta)^\alpha$ with exponent ranging from 0.8 to 1 consistent with (preliminary) predicted KBM-constrained pedestal



In NSTX, the observed width is larger than conventional tokamaks

- NSTX pedestal width is 1.7 and 2.4 larger than MAST and DIII-D & C-Mod respectively

Pedestal width scaling is consistent with predicted width for KBM constrained pedestal




- “ballooning critical pedestal”-BCP technique from EPED Model [Snyder Nucl. Fusion (2011)]
- Conventional tokamaks show an exponent of 0.5.





# Poloidal Structure Characterization using BES

---

# BES yields characterization of density fluctuations the density pedestal top





**Frequency range of interest**



D. Smith Friday Invited Y13.004


# BES provides measurements of the poloidal correlation length and poloidal velocity




- Modest change in poloidal correlation length during the inter-ELM phase
  - Poloidal correlation length corresponds to toroidal mode number ( $rk_\theta/q$ )  $n = 2 - 3$
- Measurements show ion scale fluctuation in the pedestal top

# Both BES and reflectometry systems show similar broadband power spectra

Auto-Power Spectra 45 GHz Reflectometry  
@ 665 ms (e.g., 80–99% ELM cycle)



Auto-Power Spectra BES & Reflectometry  
@ 665 ms (e.g., 80–99% ELM cycle)




Fluctuations at the pedestal top during ELM cycle are broadband as indicated by both reflectometry and BES system

# Radial Structure Characterization using Correlation Reflectometry

---

# Radial density correlation lengths at the pedestal top and steep gradient region




- The density fluctuations are measured using a 16-channel O-mode reflectometer

Crocker, PPCF (2011)

- Using two-point correlation the radial correlation function is determined.
  - tracks the equilibrium plasma motion


# 2D full wave simulation of correlation function inside pedestal region reproduces measurements



Observed correlation length corresponds to an average eddy size of  $\sim 1.3$  cm with fluctuation level in the vicinity of 1% in the gradient region.


# Reflectometry: Radial correlation length evolution depends on location inside pedestal region (steep gradient and pedestal top) UCLA

- Radial correlation length increases at the pedestal top
  - A factor of 2 increase during the last 50% of ELM cycle
  - Increase size of eddies
    - ➔ suggesting enhanced radial transport during the ELM cycle
- Steep gradient correlation length is unchanged
- Caveat: quantify the geometric effects on the measured correlation.





# Evolution of the radial displacement power spectra indicates an increase of the fluctuation level during the last 40% of ELM cycle




- Radial displacement induced by fluctuations
  - Existence of broadband fluctuations centered around 12 kHz

- Increase of the overall fluctuation amplitude late in ELM cycle
  - No clear correlation with the clamping of the pedestal gradient

- Caveat: impact of the coherent mode and 3D effects on the reflectometer signal are being investigated


# Evolution of the radial displacement power spectra indicates an increase of the fluctuation level during the last 40% of ELM cycle



- Radial displacement induced by fluctuations
  - Existence of broadband fluctuations centered around 12 kHz
- Increase of the overall fluctuation amplitude late in ELM cycle
  - No clear correlation with the clamping of the pedestal gradient
- Caveat: impact of the coherent mode and 3D effects on the reflectometer signal are being investigated

# The gyrokinetic code GENE


- GENE is a physically comprehensive Vlasov code which
  - allows for kinetic electrons electromagnetic fluctuations, collisions, and external ExB shear flows
  - is coupled to various MHD and transport codes
  - can be used as initial value or eigenvalue solver
  - supports local (flux-tube) and global (full-torus), gradient- and flux-driven simulations
  - well benchmarked and hyperscalable



*Temperature fluctuations of a global GENE simulation for ASDEX-Upgrade*

see: [gene.rzg.mpg.de](http://gene.rzg.mpg.de)  
and F. Jenko's talk

# Linear simulations using GENE at the pedestal top indicate presence of KBM-TEM hybrid modes along with microtearing modes (80 - 99% ELM)



- The characteristic scales of these instabilities appear to be consistent with experimental scales determined with BES system
- TEM modes are “hybridized” with KBM as identified by the  $\beta$  scan

# Summary/Future work

Continuous increase of the electron temperature and much less variation in electron density and pressure build up and at time saturation prior to the ELM onset

- Pressure gradient, however, is clamped during most of the ELM cycle
- NSTX exhibits wider pedestal widths than conventional tokamaks
- Pedestal width scaling like  $(\beta_\theta)^{0.8}$  in agreement with predicted KBM-constrained pedestal

Pedestal stability are performed using MHD codes

- NSTX pedestal during the last 50% of the ELM cycle is found to be kink-peeling unstable
  - Calculations were performed using ELITE code
- Using “ball” a module of GS2, pedestal top is found to be unstable to ideal high-n-ballooning modes

Characterization of the fluctuations during the inter-ELM phase

- BES and reflectometry confirm ion scale turbulence  $0.2 \leq k_\perp \rho_i \leq 0.7$
- Poloidal correlation is larger than radial correlation length

Linear GENE simulation also show the presence dominant hybrid ITG/KBM-TEM modes at the pedestal top with subdominant microtearing.

XGC1 preliminary simulation results: correlation lengths agree with experimental observations

- Most unstable mode is ITG in simulation: study will be extended to full-f nonlinear XGC1 simulation

Extend the simulation to full-f mode using XGC1 and account for measured flows and add collisions.

# Preliminary future work

---

# Simulating the Edge Turbulence during ELM cycle using XGC1

- Comprehensive gyrokinetic code
- Diverted magnetic field geometry with material wall boundary condition
  - Includes magnetic axis: wall to wall simulation
- Wall recycling of neutral particle with atomic physics
- Particle-momentum-energy conserving collision operator
- Multiscale simulation of neoclassical, **turbulence**, neutral particle, and atomic physics
  - Present XGC1 capability:
    - ITG + neoclassical + neutral in diverted geometry
    - E&M turbulence in non-diverted geometry
    - Soon to come: ITG-TEM + neoclassical + neutrals in separatrix geometry

# Nonlinear simulations using XGC1 are performed for cases during the last part of the ELM cycle

## $\delta f$ mode in XGC1

- 200 x 60 spatial grid
- simulation box up to  $\psi_n \sim 0.95$  to include the unstable region
- Using experimental profiles


Collisions and flows are not included in this simulation

- Adiabatic electrons
- Only ITG exists

Probing the fully nonlinear phase of the simulations

Characteristic poloidal structures propagating in the ion diamagnetic direction.

- ITG resides at the pedestal top, but nonlinearly and nonlocally penetrated into the pedestal region.
- Sampling a region encompassing both BES and the reflectometer measurements





# Nonlinear simulations using XGC1 are performed for cases during the last part of the ELM cycle

- $\delta f$  mode in XGC1
  - 200 x 60 spatial grid
  - simulation box up to  $\psi_n \sim 0.95$  to include the unstable region
  - Using experimental profiles
- Collisions and flows are not included in this simulation
  - Adiabatic electrons
  - Only ITG exists
- Probing the fully nonlinear phase of the simulations
- Characteristic poloidal structures propagating in the ion diamagnetic direction.
  - ITG resides at the pedestal top, but nonlinearly and nonlocally penetrated into the pedestal region.
  - Sampling a region encompassing both BES and the reflectometer measurements

# Nonlinear simulations using XGC1 are performed for cases during the last part of the ELM cycle

- $\delta f$  mode in XGC1
  - 200 x 60 spatial grid
  - simulation box up to  $\psi_n \sim 0.95$  to include the unstable region
  - Using experimental profiles
- Collisions and flows are not included in this simulation
  - Adiabatic electrons
- Probing the fully nonlinear phase of the simulations
- Characteristic poloidal structures propagating in the ion diamagnetic direction.
  - ITG resides at the pedestal top, but nonlinearly and nonlocally penetrated into the pedestal region.
  - Sampling a region encompassing both BES and the reflectometer measurements

# Nonlinear simulations using XGC1 are performed for cases during the last part of the ELM cycle

## $\delta f$ mode in XGC1

- 200 x 60 spatial grid
- simulation box up to  $\psi_n \sim 0.95$  to include the unstable region
- Using experimental profiles



Collisions and flows are not included in this simulation

- Adiabatic electrons



Probing the fully nonlinear phase of the simulations

Characteristic poloidal structures propagating in the ion diamagnetic direction.


- ITG resides at the pedestal top, but nonlinearly and nonlocally penetrated into the pedestal region.
- Sampling a region encompassing both BES and the reflectometer measurements



# Simulations from XGC1 indicate localized fluctuations with broadband power spectra



# Nonlinear simulations from XGC1 show localized fluctuations with experimental level radial and poloidal correlation lengths



# Sign-up sheet

---