


Impact of rotation and kinetic damping on the NSTX ideal-wall limit*

**Jon Menard, Zhirui Wang (PPPL)
Yueqiang Liu (CCFE)**

**APS-DPP Meeting
Providence, Rhode Island
October 31, 2012**

*This work supported by US DoE contract DE-AC02-09CH11466



*Coll of Wm & Mary
Columbia U
CompX
General Atomics
FIU
INL
Johns Hopkins U
LANL
LLNL
Lodestar
MIT
Lehigh U
Nova Photonics
Old Dominion
ORNL
PPPL
Princeton U
Purdue U
SNL
Think Tank, Inc.
UC Davis
UC Irvine
UCLA
UCSD
U Colorado
U Illinois
U Maryland
U Rochester
U Tennessee
U Tulsa
U Washington
U Wisconsin
X Science LLC*



*Culham Sci Ctr
York U
Chubu U
Fukui U
Hiroshima U
Hyogo U
Kyoto U
Kyushu U
Kyushu Tokai U
NIFS
Niigata U
U Tokyo
JAEA
Inst for Nucl Res, Kiev
Ioffe Inst
TRINITI
Chonbuk Natl U
NFRI
KAIST
POSTECH
Seoul Natl U
ASIPP
CIEMAT
FOM Inst DIFFER
ENEA, Frascati
CEA, Cadarache
IPP, Jülich
IPP, Garching
ASCR, Czech Rep*

Overview

- Drift-kinetic effects have previously been included in RWM stability analysis for NSTX (see papers by Berkery, Sabbagh, Menard)
 - In this work, we briefly review result that RWM stability is sensitive function of edge rotation, and is consistent with MARS-F analysis
 - We then investigate how RWM eigenfunctions are modified by rotation and dissipation using MARS-K
 - Not previously documented for ST plasmas
 - Could impact ‘perturbative’ approach employed by codes such as MISK
- Drift-kinetic stability analyses have ***not*** previously been carried out for ideal-wall limit, aka the ‘plasma mode’
 - Work here investigates impact of rotation, rotation shear, and kinetic damping on ‘plasma mode’ stability and eigenfunctions

Error field correction (EFC) often necessary to maintain rotation, stabilize n=1 resistive wall mode (RWM) at high β_N

- No EFC → n=1 RWM unstable
- With EFC → n=1 RWM stable



J.E. Menard et al, Nucl. Fusion 50 (2010) 045008

Analysis of experiment uses MARS: linear MHD stability code that includes toroidal rotation and drift-kinetic effects

- **Single-fluid linear MHD**

$$(\gamma + in\Omega)\xi = \mathbf{v} + (\xi \cdot \nabla\Omega)R^2 \nabla \phi$$

$$\rho(\gamma + in\Omega)\mathbf{v} = -\nabla \cdot \mathbf{p} + \mathbf{j} \times \mathbf{B} + \mathbf{J} \times \mathbf{Q} - \rho[2\Omega\hat{\mathbf{Z}} \times \mathbf{v} + (\mathbf{v} \cdot \nabla\Omega)R^2 \nabla \phi]$$

$$(\gamma + in\Omega)\mathbf{Q} = \nabla \times (\mathbf{v} \times \mathbf{B}) + (\mathbf{Q} \cdot \nabla\Omega)R^2 \nabla \phi$$

$$(\gamma + in\Omega)p = -\mathbf{v} \cdot \nabla P, \quad \mathbf{j} = \nabla \times \mathbf{Q}$$

Y.Q. Liu, et al., Phys. Plasmas 15, 112503 2008

- **Kinetic effects in perturbed p :**

$$\mathbf{p} = p\mathbf{I} + p_{\parallel}\hat{\mathbf{b}}\hat{\mathbf{b}} + p_{\perp}(\mathbf{I} - \hat{\mathbf{b}}\hat{\mathbf{b}})$$

$$p_{\parallel}e^{-i\omega t + in\phi} = \sum_{e,i} \int d\Gamma Mv_{\parallel}^2 f_L^1$$

$$p_{\perp}e^{-i\omega t + in\phi} = \sum_{e,i} \int d\Gamma \frac{1}{2} Mv_{\perp}^2 f_L^1$$

$$f_L^1 = -f_{\epsilon}^0 \epsilon_k e^{-i\omega t + in\phi} \sum X_m^u H_{ml}^u \lambda_{ml} e^{-in\tilde{\phi}(t) + im(\chi)t + i\omega_b t}$$

$$H_L = \frac{1}{\epsilon_k} [Mv_{\parallel}^2 \vec{k} \cdot \xi_{\perp} + \mu(Q_{L\parallel} + \nabla B \cdot \xi_{\perp})]$$

- **Mode-particle resonance operator:**

MARS-K:

$$\lambda_{ml} = \frac{n[\omega_{*N} + (\hat{\epsilon}_k - 3/2)\omega_{*T} + \omega_E] - \omega}{n(\langle\omega_d\rangle + \omega_E) + [\alpha(m + nq) + l]\omega_b - i\nu_{\text{eff}} - \omega}$$

MARS-F:


$$\lambda_{ml} = \frac{n[\cancel{\omega_{*N}} + (\hat{\epsilon}_k - 3/2)\cancel{\omega_{*T}} + \omega_E] - \omega}{n(\langle\cancel{\omega_d}\rangle + \omega_E) + [\alpha(m + nq) + l][\omega_b - i\nu_{\text{eff}}] - \omega}$$

+ additional approximations/simplifications in f_L^1


- **Fast ions: MARS-K: slowing-down $f(\mathbf{v})$, MARS-F: lumped with thermal**

MARS-F: Inclusion of ω_{*C} in ω_E increases separation between stable and unstable $\omega_E(\psi)$, provides consistency w/ expt.

Toroidal rotation only




Toroidal + diamagnetic




Calculated $n=1$ $\gamma \tau_{wall}$



Unstable rotation profile



Stable rotation profile





Predictions inconsistent with experiment



Predictions consistent with experiment

MARS-K studies of RWM eigenfunction: Dissipation alone can modify RWM eigenfunctions (1)



- 4% of full kinetic damping can reduce eigenfunction amplitude by 25-50% at large minor radii



NOTE: collisions are included in this and subsequent calculations with energy independent collisionality with slowing-down v evaluated at $E = 5/2$ T

MARS-K studies of RWM eigenfunction: Dissipation alone can modify RWM eigenfunctions (2)

- Full kinetic damping can produce large changes in eigenmode structure near mode rational surfaces, and in edge region




Rotation (with weak dissipation) can also modify RWM eigenfunctions

- Only small eigenfunction changes observed for $\omega_E(0)\tau_A = 0.04$
- Resonances appear on both sides of $q=2$ surface for $\omega_E(0)\tau_A = 0.1$
 - Alfvén singular points apparently split by Doppler shift



NSTX wall with $\tau_{wall}/\tau_A = 10^3$

Solid, dashed: $\alpha_D = 0.04$ $\omega_E(0)\tau_A = 0.00, 0.04$



NSTX wall with $\tau_{wall}/\tau_A = 10^3$

Solid, dashed: $\alpha_D = 0.04$ $\omega_E(0)\tau_A = 0.00, 0.10$




Shiraishi, PoP 2010


As rotation (with weak dissipation) increases toward experimental value, RWM eigenfunction is strongly modified

- Eigenfunction amplitude decreased by 25-50% at large minor radius for $\omega_E(0)\tau_A = 0.14$
- Eigenfunction strongly modified for $\omega_E(0)\tau_A = 0.16$
 - Rotation approaching marginal stability ($\omega_E(0)\tau_A \approx 0.22$)

NSTX wall with $\tau_{\text{wall}}/\tau_A = 10^3$





NSTX wall with $\tau_{\text{wall}}/\tau_A = 10^3$



Rotation + dissipation strongly modifies RWM eigenfunctions

- Rotation added to full kinetic damping produces changes that deviate significantly from cases w/o rotation or dissipation




Toroidal rotation also modifies with-wall eigenfunction

- With wall present, eigenfunction modified in both core and edge
- Note – this is ‘plasma mode’ with $\omega_r \sim$ rotation frequency

Ideal plasma, zero dissipation


Ideal wall at zero rotation marginal position

Solid: $\omega_E(0)\tau_A = 0.0$, Dashed: $\omega_E(0)\tau_A = 0.1$



Ideal wall at zero rotation marginal position

Solid: $\omega_E(0)\tau_A = 0.0$, Dashed: $\omega_E(0)\tau_A = 0.2$



Increased toroidal rotation reduces 'plasma mode' stability


- Implication: 'ideal-wall limit' is function of rotation speed
- Plasma mode predicted to be unstable for NSTX wall and rotation, but experiment does not exhibit this fast rotating instability at this time

Experimentally unstable


Ideal wall, ideal plasma, zero dissipation

Experimentally stable

Growth rate vs. rotation and wall position




Growth rate vs. rotation and wall position




Higher-resolution rotation scan finds ideal ‘plasma mode’ marginally stability at ~50-60% of experimental rotation

- **Ideal NSTX plasma: $\beta_N = 5.1$, wall position $r_{\text{wall}} / a \sim 1.25$**
 - Low rotation \rightarrow marginal $r_{\text{wall}} / a \sim 1.65$, corresponding marginal $\beta_N \sim 6$.
 - As $\omega_E(0)\tau_A \rightarrow 0.1-0.12$ (no dissipation), $n=1$ becomes unstable
 - For $\omega_E(0)\tau_A \sim 0.2-0.3$, $n=1$ mode is unstable even with the wall on the plasma boundary
- **Mode apparently mix of pressure-driven kink + Kelvin-Helmholtz**




Experimental value



With-wall limit begins decreasing from low-rotation value


Increased rotation shear in plasma core is destabilizing (consistent with expectation for Kelvin-Helmholtz)

- Rotation variations in edge region change γ very little
 - Compare **experiment** and modified **‘positive-definite’** profiles below
- γ independent of shear for **‘medium’** and **‘high’** shear cases (indicates saturation of shear effects)





**‘Positive definite’
experimental profile**

Experimental profile




However, increased rotation shear in plasma edge region can be stabilizing

- For near-edge rotation shear, both the shear magnitude and the wall position influence the mode growth rate
- ‘high’ edge-shear case is nearly stable at experimental r_{wall} / a



Plasma mode destabilization by rotation/rotation shear is still predicted when parallel sound-wave damping is included

- For larger r_{wall} / a , parallel damping systematically increases growth rate
- For smaller r_{wall} / a , growth rate can be reduced relative to ideal case



Sound wave damping model:

$$\nabla \cdot \Pi_1 = \kappa_{\parallel} \pi^{1/2} |k_{\parallel} v_{\text{thi}}| \rho \mathbf{v}_1 \cdot \mathbf{b}\mathbf{b}$$

$\kappa_{\parallel} \equiv$ Damping strength

$\kappa_{\parallel} = 1 \rightarrow$ classic ion Landau damping


$$k_{\parallel} \equiv (n-m/q)(1/R)$$

$$\text{PVISC in MARS} \equiv \kappa_{\parallel} \pi^{1/2}$$

$$\text{PVISC} = 1 \rightarrow \kappa_{\parallel} = 0.56$$

(Perpendicular) kinetic damping can stabilize 'plasma mode' over a wider range of rotation, but only at reduced r_{wall} / a


Kinetic damping from drift precession resonance included



- Mode γ generally reduced for large r_{wall} / a relative to ideal plasma predictions
- Mode can remain stable for high $\omega_E(0)\tau_A$ at small r_{wall} / a
- At sufficiently high $\omega_E(0)\tau_A \sim 0.3$, plasma unstable even for $r_{\text{wall}} / a = 1$


Results imply both rotation and dissipation influence ideal-wall stability limit (plasma mode)

Inclusion of all resonances (precession, passing, trapped) required to stabilize 'plasma mode' at experimental β , ω_E



- Including **precession** and **precession + trapped** resonances increases γ for β_N values > 4
- Inclusion of **passing particles** essential for reducing growth rate
- Stability up to $\beta_N \sim 6$ only accessed by including **all three resonance types**

Damping fraction threshold for accessing no-rotation with-wall limit of $\beta_N \sim 6$ is 30-60% of full kinetic damping



Stronger damping near experimental value provides stabilization to high β_N

Weaker damping insufficient to overcome destabilization from rotation/rotation shear

Summary

- Edge rotation ($q \geq 4$, $r/a \geq 0.8$) important for NSTX RWM
 - Trends consistent with stability calculations using MARS-F
- RWM eigenfunctions are modified by dissipation, rotation
 - Reduction/modification of ξ_{\perp} will modify kinetic stabilization
- Ideal-wall limit ('plasma mode') modified by rotation, dissipation
 - With no dissipation, plasma is predicted to be unstable at rotation $\frac{1}{2}$ the experimental value, but no instability is observed in experiment
 - Rotation shear can be stabilizing or destabilizing, depending on where the maximum shear is located in minor radius
 - Parallel (SW) damping destabilizing/stabilizing at large/small r_{wall} / a
 - Inclusion of full kinetic damping stabilizes plasma mode at high rotation, β
 - Passing resonances appear most important for plasma mode stability
 - Next-steps: Compare experimental mode frequencies and eigenfunctions to predicted values including rotation and full kinetic resonant effects