


Supported by




# Internal Transport Barriers in NSTX Plasmas? – A Progress Report

Martin Peng, Ron Bell, Ben LeBlanc, Ed Synakowski,  
Jon Menard, Dave Gates, Stan Kaye,  
Steve Sabbagh, Dan Stutman

for the NSTX National Team

**5<sup>th</sup> Meeting of the ITPA  
Transport & ITB Physics Topical Group**

29 September – 2 October, 2003



Columbia U  
Comp-X  
General Atomics  
INEL  
Johns Hopkins U  
LANL  
LLNL  
Lodestar  
MIT  
Nova Photonics  
NYU  
ORNL  
PPPL  
PSI  
SNL  
UC Davis  
UC Irvine  
UCLA  
UCSD  
U Maryland  
U New Mexico  
U Rochester  
U Washington  
U Wisconsin  
Culham Sci Ctr  
Hiroshima U  
HIST  
Kyushu Tokai U  
Niigata U  
Tsukuba U  
U Tokyo  
JAERI  
Ioffe Inst  
TRINITY  
KBSI  
KAIST  
ENEA, Frascati  
CEA, Cadarache  
IPP, Jülich  
IPP, Garching  
U Quebec

# This Study Examines NSTX Data for the Near-Term Tasks Defined by ITPA Topical Group on ITB&T



## In 1-2 Years:

- Improve experimental understanding of critical issues of burning plasmas with ITB:  $T_i \sim T_e$ , low  $V_\phi$ , flat density profile,  $Z_{\text{eff}} < 2$ 
  - ***ITB formation, evolution, and sustainment conditions***
  - ***Impurity accumulation***
  - ***Compatibility with divertor requirements***
- Develop, manage, and analyze new experimental ITB database
- Test simulation and modeling of ion transport
  - ***e.g., JT-60U “box-like” ITB  $T_i$  profiles, JET  $(r/a)_{\text{ITB-foot}}$  evolution, etc.***

**NSTX Data Are Being Compiled and Analyzed to Address Issues in:**

- **ITB formation, evolution, sustainment**
- **Impurity accumulation**

# NSTX Has Built up Basic and Modern Diagnostic Capabilities to Support Research



## Core Plasma Diagnostics

- Thomson scattering (20 ch., 60Hz)
- Charge Exchange Recomb. Spect. (CHERS):  $T_i$  &  $v_\phi$  (51 ch.)
- VB detector (single chord)
- Soft x-ray arrays (4) [JHU]
- Bolometer array (midplane tangential)
- X-ray crystal spectrometer ( $T_i(0)$ ,  $T_e(0)$ )
- Edge rotation spectroscopy
- Electron Bernstein wave radiometer
- FReTIP interf'r/polarim'r (4 ch) [UCD]
- PICXIS Fast 2D X-ray camera [Frascati, JHU]
- Tang. X-ray pin hole camera [U. Wisconsin]

## Magnetics and MHD

- Magnetics for equilibrium reconstruction
- Diamagnetic flux measurement
- High-n and high-frequency Mirnov arrays
- Locked mode coils
- 1mm interferometer [UCLA]

## Turbulence

- Edge reflectometer [UCLA]
- Edge fluctuation imaging [LANL, PSI]

## Plasma Monitoring

- Fast visible camera [LANL]
- VIPS: Visible spectrometer
- SPRED: UV spectrometer
- Transmission grating spectrometer [JHU]
- EFIT (Columbia University)



## Boundary Physics

- Divertor Bolometer
- Fast probe [UCSD]
- Infrared Camera (2) [ORNL]
- Fast Ion Gauge [University of Wash]
- Divertor fast camera [Hiroshima Univ.]
- Divertor tile Langmuire probe array
- 1-D CCD  $H_\alpha$  camera (2) [ORNL]
- Visible filterscopes ( $H_\alpha$ , OII, CII) [ORNL]
- Scrape-off layer reflectometer [ORNL]
- Fast camera (PSI)

## Energetic Particles

- Fission chamber neutron measurement
- Fast neutron measurement
- Neutral particle analyzer (scanning)
- Fast ion loss probe





# Co-NBI, High- $\beta_p$ ( $\sim 1$ ), Sustained ELM-Free H-Mode Plasmas Are Appropriate for ITB Studies



$$f_{BS} \sim 0.5; f_{NBI} \sim 0.1; V_L \sim 0.1 \text{ V for } \geq \tau_{Skin}$$

Gates, Menard, Sabbagh

# $T_i$ Substantially Higher Than $T_e$ in Plasma Core for the “Flattop” Duration



20 40 60 80 100 120 140 160



20 40 60 80 100 120 140 160

RADIUS (cm)

RADIUS (cm)

# Recent EPS Poster on ITB Provided Definition for ITB Formation




- “ITB foot” is located by **peak of  $(d/dR)(R_0/L_{Ti})$** ,  $\sim$  **peak of  $dL_{Ti}/dR$**
- Critical values are defined for the “ITB foot”
  - $R_0/L_{Ti}$  &  $\rho_i^* = \rho_i/L_{Ti}$
- Also value of  $\rho_i/R_0$  at “ITB foot”




# NSTX Routinely Exhibits Similar Behavior Under NBI Just Before H-Mode Transition





- >20 ms before H-mode transition after 2 NBI source power
- Peak ( $dL_{Ti}/dR$ ) is clearly located
- Critical values measured
  - $R_0/L_{Ti}$  &  $\rho_i^* = \rho_i/L_{Ti}$
- And value of  $\rho_i/R_0$



# NSTX Data Generally Near or Beyond the Boundary of the Tokamak Range, Showing High Leverage





“It is suggested that critical values of  $\rho_i/L_{Ti}$  depends on other quantities than  $\rho_i$  and  $L_{Ti}$ ” – Fujita





# The Peak ( $dL_{Ti}/dR$ ) Location Moves to Plasma Edge in ~50 ms After H-Mode Transition





Need 51-ch. CHERS (ready), and MSE to estimate  $q(R)$ .


Plasma Edge

R. Bell, B. LeBlanc

# Location of Steep $T_i$ & $V_\phi$ Gradients Broadens and Moves Outward Following H-Mode Transition



# Hollow Zeff (Carbon) Profile Maintained After H-mode Formation, Similar to VH-Mode?




R. Bell, B. LeBlanc

# Gyrokinetic Microinstability Calculations Indicate Suppression of Weak ITG by Flow Shear





- NBI-driven flow shearing rate  $\gg$  ITG growth rate ( $T_i \sim 2T_e$ )
- Virulence of ETG depends strongly on  $T_i/T_e$ 
  - not likely stabilized by flow shearing for  $T_e \leq T_i$
- Other physics under exploration
  - effects of  $\beta'$
  - stabilization by negative magnetic shear



## Gyrokinetic Microinstability Growth Rates



# Location of Minimum $\chi_i$ Moves Outward as Beam Ion Heating Profile Remains Broad for $R < 1.4\text{m}$



# 51 Chord CHERS Data of 2003 Shots Will Shed More Light on Detail of ITBs in NSTX



• “Flat spot” in  $F_\phi$  appears to be associated with 2/1 MHD island

# ITB-Relevant Data from NSTX Are Intriguing and Been Analyzed for ITPA and ST Research



- $T_i > T_e$  over substantial regions; microinstability analysis suggests suppressed ITG modes.
- $T_i$  gradient steepens over small region before H-mode transition (not  $T_e$ ); preliminary analysis suggests local minimum in  $\chi_i$ .
- The regions of peak barrier  $(dL_{Ti}/dR)_{peak}$  and  $(dV_\phi/dR)_{peak}$  move toward edge region within 50 ms of H-mode transition; what mechanisms drive this evolution?
- Hollow Zeff (carbon) profile maintained after H-mode formation; is this similar to VH-mode in standard A?
- Started neoclassical momentum balance calculations; strong toroidal flow ( $V_\phi/V_{Alfven} \leq 0.3$ ) effects?
- High resolution CHERS, edge flow spectroscopy, MSE; microinstability calculations, ExB shear profile?
- Much work to be done and for comparison with standard A