


# Impurity transport in electron-dominated RF-heated scenarios

L. Delgado-Aparicio<sup>1</sup>, J. Rice<sup>2</sup>, K. Tritz<sup>3</sup>, J. Muñoz<sup>3</sup>,  
D. Stutman<sup>3</sup>, D. Smith<sup>4</sup>, N. Bertelli<sup>1</sup>, R. Perkins<sup>1</sup>,  
J. Hosea<sup>1</sup>, R. E. Bell<sup>1</sup>, F. Poli<sup>1</sup>, H. Yuh<sup>5</sup>, A. Diallo<sup>1</sup>,  
B. Leblanc<sup>1</sup>, W. Wang<sup>1</sup> and W. Gutenfelder<sup>1</sup>

<sup>1</sup>PPPL, <sup>2</sup>MIT Plasma Science and Fusion Center, <sup>3</sup>JHU,  
<sup>4</sup>University of Wisconsin-Madison, <sup>5</sup>Nova Photonics

NSTX-U Research Forum, B318, Feb. 24-27<sup>th</sup>, 2015



Coll of Wm & Mary  
 Columbia U  
 CompX  
 General Atomics  
 FIU  
 INL  
 Johns Hopkins U  
 LANL  
 LLNL  
 Lodestar  
 MIT  
 Lehigh U  
 Nova Photonics  
 ORNL  
 PPPL  
 Princeton U  
 Purdue U  
 SNL  
 Think Tank, Inc.  
 UC Davis  
 UC Irvine  
 UCLA  
 UCSD  
 U Colorado  
 U Illinois  
 U Maryland  
 U Rochester  
 U Tennessee  
 U Tulsa  
 U Washington  
 U Wisconsin  
 X Science LLC


Culham Sci Ctr  
 York U  
 Chubu U  
 Fukui U  
 Hiroshima U  
 Hyogo U  
 Kyoto U  
 Kyushu U  
 Kyushu Tokai U  
 NIFS  
 Niigata U  
 U Tokyo  
 JAEA  
 Inst for Nucl Res, Kiev  
 Ioffe Inst  
 TRINITI  
 Chonbuk Natl U  
 NFRI  
 KAIST  
 POSTECH  
 Seoul Natl U  
 ASIIPP  
 CIEMAT  
 FOM Inst DIFFER  
 ENEA, Frascati  
 CEA, Cadarache  
 IPP, Jülich  
 IPP, Garching  
 ASCR, Czech Rep

# This proposal aims to characterize impurity transport in electron-dominated RF-heated scenarios

## FACTS

- a) In ASDEX H-modes with dominant auxiliary ion heating by NBI the impurity particle pinch is regularly measured directed inwards.
- b) In TCV L-modes with dominant auxiliary  $e^-$  heating by ECRH, the impurity pinch is measured directed outwards.
- c) In JET H-modes using NBI + ICRH, the pinch velocity has been systematically found to change its direction from strongly inwards to slightly outwards when using ICRH in MH (ions) vs MC (electrons).
- d) New experiments @ C-MOD!

$\varepsilon_{\text{SXR,Ca}}$  (Ohmic vs LHCD – AFTER Ca inj.)



L. Delgado-Aparicio and J. Rice, MP# 710  
Alcator C-Mod @ MIT-PSFC  
(work in progress to support JRT15)

# Is turbulent impurity transport in play? TEM modes are leading candidate for “explain” core shielding

## a) TEP pinch:

- a.k.a: B curvature pinch

**TEM (-) ITG (-)**

$$\frac{V_Z^{TEP}}{D} \approx -\frac{4s/3 + 0.5}{R}$$

## b) Thermodiffusion pinch:

- Depend on the  $v_{\text{phase}}$ .


**TEM (-) ITG (+)**

$$\frac{V_Z^{TD}}{D} \approx \frac{1}{Z} \cdot \frac{\nabla T_Z}{T_Z}$$

## c) Parallel compressibility:

- $\nabla T$  driven
- the  $T_e/T_i$  dependence is negligible
- $\propto Z/m_Z$
- mass/charge independent for fully stripped impurities:  
e.g.  $C \Rightarrow Z_C/m_C = 1.0/m_D$ ,  $Ne \Rightarrow Z_{Ne}/m_{Ne} = 1.0/m_D$
- Predict the effect for Moly:  $Mo \Rightarrow Z_{Mo}/m_{Mo} = 0.67/m_D$

**TEM (+) ITG (-)**



# 1-day experiment can help elucidate role of turbulent particle transport in NSTX-U

- Impact operation with high-Z PFCs.
- Probes particle transport in the **absence of particle and momentum sources!**
- Has NEVER been tested in NSTX.
- This is a particular attractive scenario for **ITER** since  $\alpha$ -heating predominantly heats the electron channel.
- Will provide valuable data for **NSTX-U T&T thrusts #s 2 & 3**, and **ITPA TC #s 11 & 15**.
- Request **one day of experiments** w/ and w/o **Ne-puffs** while scanning the **HHFW power** from 0.5 to 3 MW (no-Lithium). Key diagnostics are CHERS, MPTS, MSE and the new ME-SXR [5] and AXUV-based “bolometer” [6] systems.
- Key diagnostics are: CHERS, MPTS, MSE, USXR and the new ME-SXR and AXUV-based “bolometer” systems.