


Impurity transport vs torque in NBI heated H-Modes

L. Delgado-Aparicio¹, K. Tritz², J. Muñoz², J. Rice³,
D. Stutman², D. Smith⁴, R. E. Bell¹, F. Poli¹, H. Yuh⁵,
A. Diallo¹, B. Leblanc¹, W. Wang¹ and W. Gutenfelder¹

¹PPPL, ²JHU, ³MIT Plasma Science and Fusion Center,

⁴University of Wisconsin-Madison, ⁵Nova Photonics

NSTX-U Research Forum, B318, Feb. 24-27th, 2015




Coll of Wm & Mary
 Columbia U
 CompX
 General Atomics
 FIU
 INL
 Johns Hopkins U
 LANL
 LLNL
 Lodestar
 MIT
 Lehigh U
 Nova Photonics
 ORNL
 PPPL
 Princeton U
 Purdue U
 SNL
 Think Tank, Inc.
 UC Davis
 UC Irvine
 UCLA
 UCSD
 U Colorado
 U Illinois
 U Maryland
 U Rochester
 U Tennessee
 U Tulsa
 U Washington
 U Wisconsin
 X Science LLC

Culham Sci Ctr
 York U
 Chubu U
 Fukui U
 Hiroshima U
 Hyogo U
 Kyoto U
 Kyushu U
 Kyushu Tokai U
 NIFS
 Niigata U
 U Tokyo
 JAEA
 Inst for Nucl Res, Kiev
 Ioffe Inst
 TRINITY
 Chonbuk Natl U
 NFRI
 KAIST
 POSTECH
 Seoul Natl U
 ASIPP
 CIEMAT
 FOM Inst DIFFER
 ENEA, Frascati
 CEA, Cadarache
 IPP, Jülich
 IPP, Garching
 ASCR, Czech Rep

This proposal aims to characterize impurity transport with the 2nd NBI (off-axis heating and torque)

- Other expts. will revisit the Z-transport ρ^* and v^* experiments performed with NBI #1 in NSTX (see J. Muñoz *et al.*).
- Goal is to characterize the Z-transport properties of H-mode scenarios with high toroidal velocities using the 2nd NBI.
- Off-axis beams may bring a concomitant increase of the D & Z Mach numbers providing additional momentum to the outward low temperature plasma.
- The presence of poloidal asymmetries and off-axis impurity peaking is nearly guaranteed.
- Mach correction is proportional to the m_z so the effects impurities will be easily distinguishable from that of D or C.



Synthetic $P_{\text{rad,Fe}}$ for $P_{\text{rad,Tot}} = 1$ MW.
(L. Delgado-Aparicio, RSI'14)

Off-axis NBI torque can change Z-Mach numbers leading to off-axis impurity peaking

- Understanding the radial dependence of poloidal asymmetries and its role for the outward “convection” of impurities is highly desirable.
- Impact operation with high-Z PFCs.
- Will provide valuable data for the **JRT2015** (“Quantify impact of broadened J & p profiles on confinement and stability”) and **NSTX-U T&T thrusts #s 2 & 3**, and **ITPA TC #s 11 & 15**.
- Request **one day of experiments** w/ and w/o **Ne-puffs** and changing the **off-axis torque density** ($R_{tg} \sim 110, 120$ and 130 cm) using the expanded NBI flexibility (e.g. at 4, 5 or 6 MW). No-Lithium.
- Key diagnostics are: **CHERS**, **MPTS**, **MSE**, **USXR** and the new **ME-SXR** and **AXUV**-based “bolometer” systems.

